



Bunions are a **progressive disorder** and do not go away on their own without surgical correction.¹

About
1 in 4
US adults
suffer from
bunions.²



Up to 30% of patients
with bunions also have MTA.^{3,4}



Adductoplasty®
Midfoot Correction System

Lapiplasty®
3D Bunion Correction

Individual results may vary depending on age, weight, health and other variables.

There are risks and recovery takes time.

Risks include infection, pain, implant loosening and loss of correction with improper bone healing. If any of these occur, additional treatments may be needed.

For more information on benefits, risks and recovery, visit Lapiplasty.com.

1. American College of Foot and Ankle Surgeons (ACFAS) Website ©2022

2. Nix S, et al. J Foot Ankle Res. 2010. 27:3:21.

3. Aiyer AA, et al. FAJ 2016; 37: 165-171.

4. Aiyer AA, et al. FAJ. 2014; 35: 1292-1297.

TREACE
Medical Concepts, Inc.
The Leader in Hallux Valgus Surgery™

Registered marks and trademarks of Treace Medical Concepts, Inc.
[Pat. treace.com/patents](http://Pat.treace.com/patents)

©2024 Treace Medical Concepts, Inc. All Rights Reserved. M2104B

Adductoplasty®
Midfoot Correction System

**Because not all
bunions are the same.**




Complementing Lapiplasty® 3D Bunion Correction.

Know Your Bunions and Your Options

What is Metatarsus Adductus (MTA)?


MTA is a common foot deformity that causes the front half of the foot, or forefoot, to turn inward. A full correction of the bunion may require repositioning the bones of the 2nd and 3rd toes in order to make room for complete first toe realignment. Failure to address MTA increases the risk of bunion recurrence.³



Midfoot Malalignment
Blocks Bunion Correction

Why have Adductoplasty?

The **Adductoplasty® Procedure** corrects the alignment of the 2nd and 3rd toes, creating the necessary space to perform the **Lapiplasty® Procedure** and bring the big toe back into normal alignment.



Midfoot Alignment
Allows Full Bunion Correction

The Lapiplasty® Family of Procedures

Not all bunions are the same. Sometimes bunions are accompanied by big toe arthritis or a midfoot deformity like MTA. Adductoplasty® instrumentation, specifically designed to correct this deformity, along with the associated implants, provide a comprehensive system that allows for realignment and stabilization of the midfoot.



Bunion without MTA



Correction with Lapiplasty®
3D Bunion Correction



Bunion with MTA



Correction with
Adductoplasty® and
Lapiplasty® Procedure